

SEDAP

A PROGRAM FOR RESEARCH ON

SOCIAL AND ECONOMIC DIMENSIONS OF AN AGING POPULATION

**Agreement between Self-Reported and
Routinely Collected Health Care Utilisation
Data among Seniors**

**Parminder Raina, Vicki Torrance-Rynard
Micheline Wong, Christel Woodward**

SEDAP Research Paper No. 81

For further information about SEDAP and other papers in this series, see our web site:
<http://socserv2.mcmaster.ca/sedap>

Requests for further information may be addressed to:
Secretary, SEDAP Research Program
Kenneth Taylor Hall, Room 426
McMaster University
Hamilton, Ontario, Canada
L8S 4M4
FAX: 905 521 8232
e-mail: qsep@mcmaster.ca

**Agreement between Self-Reported and Routinely
Collected Health Care Utilisation Data among Seniors**

**Parminder Raina, Vicki Torrance-Rynard
Micheline Wong, Christel Woodward**

SEDAP Research Paper No. 81

September 2002

The Program for Research on Social and Economic Dimensions of an Aging Population (SEDAP) is an interdisciplinary research program centred at McMaster University with participants at the University of British Columbia, Queen's University, Université de Montréal, and the University of Toronto. It has support from the Social Sciences and Humanities Research Council of Canada under the Major Collaborative Research Initiatives Program, and further support from Statistics Canada, the Canadian Institute for Health Information, and participating universities. The SEDAP Research Paper series provides a vehicle for distributing the results of studies undertaken by those associated with the program. Authors take full responsibility for all expressions of opinion.

Agreement between Self-Reported and Routinely Collected Health Care Utilisation Data among Seniors

Parminder Raina, PhD., Vicki Torrance-Rynard, MSc., Micheline Wong, MHSc., Christel Woodward, PhD.

Parminder Raina, Ph.D., is Associate Professor in the Department of Clinical Epidemiology and Biostatistics, McMaster University. He was formerly Assistant Professor of Epidemiology in the Department of Health Care and Epidemiology, Faculty of Medicine, University of British Columbia, and the Centre for Community Health and Health Evaluation Research, where he was Director of the BC Injury Research and Prevention Unit. Dr. Raina is a recipient of Health Canada's National Health Scholar Award.

Vicki Torrance-Rynard, MSc., is a Research Coordinator in the Department of Clinical Epidemiology and Biostatistics, Faculty of Health Sciences, McMaster University, Hamilton, Ontario Canada.

Micheline Wong, MHSc., is a Research Assistant at the Department of Health Care and Epidemiology, Faculty of Medicine, University of British Columbia, and the Centre for Community Health and Health Evaluation Research in Vancouver, British Columbia, Canada.

Christel Woodward, Ph.D., is a Professor in the Department of Clinical Epidemiology and Biostatistics, Faculty of Health Sciences, McMaster University, and the Centre for Health Economics and Policy Analysis, Faculty of Health Sciences, McMaster University, Hamilton, Ontario, Canada

Correspondence:

Parminder Raina, Associate Professor
McMaster University
Evidence Based Practice Centre
Courthouse T-27 Building, 3rd Floor
1280 Main Street West
Hamilton, Ontario, CA L8S 4L8
Ph: 905-525-9140, Ext. 22197
Fax: 905-522-7681
e-mail: praina@mcmaster.ca

Abstract

Objective: To examine the agreement between self-reported and routinely collected administrative health care utilisation data, and the factors associated with agreement between these two data sources.

Data Sources/Study Setting: A representative sample of seniors living in an Ontario county within Canada was identified using the Ontario Ministry of Health's Registered Persons Data Base in 1992. Health professional billing information and hospitalisation data were obtained from the Ontario Ministry of Health and Long-Term Care (OMH) and the Ontario Health Insurance Plan (OHIP).

Study Design: A cross-sectional survey was carried out to assess *any* contact and *frequency* of contacts with health professionals and hospital admissions. Similar information was obtained from routinely collected administrative data. The level of agreement was assessed using the proportion of absolute agreement, Cohen's kappa statistic (κ), and the intraclass correlation coefficient (ICC). Logistic and linear regression were used to identify factors that were associated with the magnitude and direction of disagreement respectively.

Data Collection/Extraction Methods: Telephone interviews were conducted on 1,054 seniors, and completed data were available for 1,038 seniors. Each respondent's personal health number was used to electronically link survey data with health professional billing and hospitalisation data bases.

Principal Findings: Substantial to almost perfect agreement was found for the contact utilisation measures, while agreement on volume utilisation measures varied from poor to almost perfect. In surveys, seniors overreported contact with general practitioner and physiotherapists or

chiropractors, and underreported contact with other medical specialists. Seniors also underreported the number of contacts with general practitioners and other medical specialists. The odds of agreement decreased if respondents were male, aged 75 years and older, had incomes of less than \$25,000, had poor/fair/good self-assessed health status, or had two or more chronic conditions.

Conclusion: The findings of this study indicate that there are substantial discrepancies between self-reported and administrative data among older adults. Researchers seeking to examine health care use among older adults need to consider these discrepancies in the interpretation of their results. Failure to recognize these discrepancies between survey and administrative data among older adults may lead to the establishment of inappropriate health care policies.

Keywords: health services utilisation, seniors, self-reports, agreement, billing data

Introduction

Epidemiological and health services research has increasingly focused attention on health care utilisation among the aging population (Ory and Bond 1989). Many of these studies have consistently shown strong associations between health problems, aging and the use of health care services. However, it remains to be seen whether improved health practices and changes to health care delivery will translate into better individual health for seniors in the future. The increasing numbers of seniors in many industrialised countries has already placed a growing demand on the health care system (Mossey, Havens and Wolinsky 1989). In recent years, health services research has closely examined the development of accurate measures of health care utilisation by seniors, as well as factors affecting the use of health care services (Glandon, Counte and Tancredi, 1992; Wan 1989; Muller 1986). In many of these studies, self-reported questionnaires have been the most common method of collecting information on respondents' health status and their use of health care services. Roos and Havens (1991) found that self-reported health was a strong predictor of "successful aging" or remaining independent, and that those who aged successfully made markedly fewer demands on the health care system. Other studies in the literature have also found a positive association between self-reported health and the use of health care services among the elderly population (Prigerson, Maciejewski, Rosenheck 1999; Raina 1995; Branch, Jette and Evashwick 1981).

Self-reported questionnaires, however, rely on respondents' ability to recall past events. It is important to understand the accuracy of survey responses among seniors, especially since seniors have been known to, both over and underreport their use of health care services more often than younger individuals (Cummings and Nevitt 1988; National Center for Health Statistics 1987). The National Center for Health Statistics in the United States examined the accuracy of

self-reported utilisation and found significant underreporting of hospitalisation days by respondents aged 55 years and older compared with those under the age of 55 (National Center for Health Statistics 1987). Studies specifically on seniors have found substantial over- and underreporting of health care services (Cummings, Nevitt and Kidd 1988; Bush, Miller, Golden et al. 1989; Mackenbach, Looman and Van der Meer 1996; Wallihan, Stump and Callahan 1999). Wallihan and colleagues (1999), for example, studied a group of 4,506 patients aged 60 years and older. They found that approximately one-quarter of older adults failed to report a hospitalisation in the past 12 months and almost one-half underestimated the number of hospitalisations by at least one episode.

The ability or willingness to accurately report use of health care services may decline with physical and cognitive function (National Center for Health Statistics 1987; Wallihan, Stump and Callahan 1999). The National Center for Health Statistics found that more acute and chronic health conditions, and longer length of stay in the hospital were related to less underreporting (National Center for Health Statistics 1987). The possibility that discrepancies exist has important implications for future delivery and accessibility of health care services, particularly if survey findings play an important role in policy-making decisions. If the extent of over- and underreporting of health care utilisation do not balance with each other, the average utilisation of services in the elderly population may be much different than is currently believed (Glandon, Counte and Tancredi 1992). Moreover, the impact that inaccurate findings have on more complicated statistical analyses relating to health utilisation is likely to be even more dramatic. For instance, if older and sicker individuals tend to over-report the use of services systematically, then the empirical findings from studies will tend to estimate a greater effect of age and health status on health utilisation than actually exists. These reasons emphasize the importance of

studies to assess the factors that affect reported health care utilisation. The differences observed between self-reported and routinely collected health utilisation data in community-based seniors population are not well documented in the literature (Glandon, Counte and Tancredi 1992). Few studies, if any, have attempted to identify factors that are associated with the level of agreement observed.

This study, used a representative sample of community-based non-institutionalised seniors to study the level of agreement between self-reported and routinely collected health care utilisation data. In addition, this study explored salient factors that may be associated with agreement or disagreement between the two data sources.

Methods

Study sample

The target population for this study was community-dwelling adults aged 65 years and older. A sample of (n=1500) of older people stratified by age and sex was identified using the Ontario Ministry of Health's (MOH) Registered Persons Data Base (RPDB). The RPDB is an ideal sampling frame for selecting a random sample of subjects because of Canada's universal healthcare system. The RPDB contains demographic information such as names, date of birth, sex, and address for all residents registered for healthcare coverage in Ontario.

From the sample of 1,500 seniors, 1,296 consented to participate in the study. However, when contacted for the telephone interview, only 1,054 agreed to participate. Complete data for the study were available for 1,038 seniors. Comparison between respondents and non-respondents showed that both groups were similar for factors such as gender, marital status, education, physical activity and perceived health status. However, respondents were significantly more likely to be younger in age and have a higher household income than non-respondents

(Raina, Bonnett, Waltner-Toews, et al. 1999). In November 1992, trained interviewers conducted telephone interviews approximately 30 minutes in length. Further description of the study and survey design is available elsewhere (Raina, Waltner-Toews, Bonnett et al. 1999).

Questionnaire

The questionnaire used in the study assessed two types of health care utilisation, contacts with health professionals and hospital admissions. For contacts with health professionals, respondents were asked, “*During the past 12 months, have you had contact with any of the following about your physical or mental health: a) general practitioner or family physician? b) medical specialist (such as a cardiologist, urologist or psychiatrist)? c) physiotherapist or chiropractor?*”. If a respondent answered “yes”, they were then asked about the number of times they saw that particular health professional. For hospital admissions, respondents were asked, “*Were you in a hospital overnight during the past 12 months?*”. If a respondent answered “yes” they were then asked for the number of nights they spent in the hospital. These questions were adopted from the Statistics Canada’s 1994 National Population Health Survey (NPHS) (Statistics Canada 1994). A 12-month recall period for each question on health care was used to compare the results with national surveys. The test-retest reliabilities for these questions were found to be moderate to high; interclass correlation coefficient (ICC) ranged from 0.69 to 0.91 (Prigerson, Maciejewski and Rosenheck 1999).

The questionnaire also asked about a variety of variables that have been shown in the literature to be associated with health care utilisation (Wan 1989; Roos and Havens 1999; Chappell, Strain and Blandford 1986). The following socio-demographic variables were included: gender (female or male); age (i.e. 65-69 years old, 70-74 years old, 75-79 years old, or 80 years and over); gross household income level (under \$11,999, \$12,000 to \$24,999, \$25,000

to \$49,999, or \$50,000 and over); education (less than high school graduation, or high school and above); current marital status (married, widowed or other which included never married, divorced and separated); living arrangement (alone or living with someone); and type of dwelling (house, apartment or other which included trailer, townhouse and other types of dwellings). In addition, current status of smoking cigarettes (current smokers, past smokers or never smokers) and alcohol drinking (never drinkers, sometimes drinkers or regular) were also assessed. For the category of current status of alcohol drinking, “never drinkers” were respondents who did not drink alcohol at all, “sometime drinkers” drank 1 to 3 times a month or less than once a month, and “regular drinkers” drank once a week or more (Raina 1995).

Health status variables in the study included self-perceived health status, self-perceived emotional health status, and number of chronic health conditions. Self-perceived health status was assessed by asking the question, “*In general, compared to other persons your age, would you say your health is...excellent, very good, good, fair or poor?*” Self-perceived emotional health status was based on the question, “*How would you rate your mental or emotional health at the present time... excellent, very good, good, fair, poor or very poor?*” The number of chronic conditions experienced by respondents were assessed by answering “yes” or “no” to whether a doctor had ever diagnosed the respondent with any of the following health problems: arthritis, asthma, back pain, chronic bronchitis, diabetes, epilepsy, heart trouble, cancer, high blood pressure, stroke, cataract, glaucoma, or stomach/intestinal ulcers. Respondents were then classified as having zero, one, two, or three or more chronic illnesses. These questions were also adopted from the national surveys and were shown to have moderate to high test-retest reliabilities. The interclass correlations (ICC) ranged from 0.69 to 0.96 (Raina, Bonnett, Waltner-Toews, et al. 1999; Prigerson, Maciejewski and Rosenheck 1999;).

Ministry of Health and Long-Term Care Utilisation Data

This study utilised two data bases of routinely collected administrative data from the Ontario Ministry of Health and Long-Term Care (OMH) and the Ontario Health Insurance Plan (OHIP): health professional billing information and hospitalisations. Respondents provided written permission to access their health care data from the OMH. As part of the public-funded health care system in Canada, health care utilisation data on both outpatient and inpatient services are routinely maintained by each province. Residents who are covered by the health care system are issued a unique personal health number (PHN) which entitles them to health care services paid for by the province's health insurance plan. OHIP maintains a health professional billing data base on the specific services paid to health professionals when they submit a claim for services rendered. This data base includes information on the following: the patient's personal health number, provider identification code, type of specialty, diagnostic and procedure codes, referring physician, date of service, fee schedule code, number of services and fee(s) paid.

Similarly, the hospitalisation data base, maintained by the OMH, records information on all inpatient admissions to any Ontario hospital, including personal health number, date of admission, date of discharge, diagnostic and procedure codes and length of stay. In this study, each respondent's PHN was used to electronically link his/her survey data with the OHIP billing and hospitalisation data base. Information from billing and hospital data bases were extracted for the year immediately preceding the survey data (November 1, 1991 to October 31, 1992) to ensure that the time frame associated with all sources of data coincided. For the purposes of the study, we identified physicians by their area of specialty rather than the specific services that were billed to OHIP. To verify the physician's area of specialty, an OHIP demographic data base containing information on the original specialty declared by the health professional and other

related data, such as age, gender, place of graduation, year of graduation, type of specialty and provider identification code, was used in the study. Discrepancies between the billing data base and the demographic data base, however, occurred in less than 2% of listed clinicians.

Analysis

Descriptive statistics were calculated for all independent variables available in the study. The level of agreement for any stay in a hospital and any contact with a health professional, as well as for the number of nights stayed in hospital and the number of contacts with a health professional were examined. Ten records from the health professional utilisation data base and two records from the hospitalisation data base were dropped due to missing data leaving a final sample size of 1,028 respondents in the health professional analyses and 1,036 in the hospital analyses. For the purposes of this study, the measures of any stay in hospital (yes or no) and any contact with a health professional (yes or no) were termed “contact utilisation measures”. The measures related to the number of nights stayed in hospital and the number of contacts with a health professional were termed “volume utilisation measures”.

The level of agreement for dichotomous contact utilisation was assessed using the proportion of absolute agreement and beyond chance agreement using Cohen’s kappa statistic (κ). Agreement for continuous volume utilisation measures was assessed using the random effects intraclass correlation coefficient (ICC) (Strout and Fleiss 1979; Fleiss 1986). The 95% confidence intervals around the κ and ICC were calculated (Strout and Fleiss 1979; Fleiss 1986). The benchmark for determining the closeness of the comparison for both κ and ICC was based on Landis and Koch (1977). In their scale, the strength of agreement was as follows: 0.00-0.20 (poor), 0.21-0.40 (fair), 0.41-0.60 (moderate), 0.61-0.80 (substantial), and 0.81-1.00 (almost perfect).

Agreement for the contact utilisation measures included “yes/yes” and “no/no” responses in the self-reported and administrative data respectively. Disagreement, therefore, included “yes/no” (overreporting in survey) and “no/yes” (underreporting in survey) responses. The volume utilization measures were also analysed in a similar manner.

Polytomous logistic regression for categorical outcomes would have been the most appropriate to assess the predictors of under or overreporting for the contact utilisation measures. However, due to small sample size, each dependent variable for the contact utilisation measures was dichotomised. Each dependent variable was coded as “1” indicating agreement between the two data sources or “0” indicating disagreement (included over and underreporting). Each independent variable known to be associated with health care utilisation was selected and used in the logistic regression to identify predictors of agreement/disagreement for contact utilisation measures. Separate logistic regressions were run for the following four dependent variables: any stay in hospital, any contact with GP (general practitioner), any contact with physiotherapist (PT) or chiropractor, and any contact with other medical specialties. Each variable used in the study was examined for its effect on each model. Variables were omitted from the analyses if they were not significantly associated with agreement in health care utilisation measures ($p > 0.05$) and did not contribute to the overall model. The variables of age and income level were collapsed into smaller categories due to small sample sizes. Age was regrouped into two categories, 65-74 years old and 75 years and over. Household income was regrouped according to whether individuals earned \$25,000 or less, or more than \$25,000.

For volume utilisation measures, linear regression was used to assess the predictors of under or overreporting. The dependent variable was the difference between self-reported and MOH data. Positive values imply that self-reported measures of utilization exceed the measures

of utilisation in the MOH data (e.g., overreport). Negative values imply that self-reported measures of utilisation are less than the measures of utilisation in the MOH data (e.g., underreport). The same independent variables used in the above mentioned logistic regression appear in each of the four difference linear regression equations. In addition, in each of the four difference equations (e.g., difference in length of stay), the respective volume utilisation measure from the MOH data was included as a independent variable (e.g., length stay from MOH data). Inclusion of volume measure utilisation from the MOH data as an independent variable controlled for estimation bias (Glandon, Counte and Tancredi 1992).

Results

Demographic Characteristics

The sample size consisted of 1,038 seniors with a mean age of 73 years ($SD=\pm 6.3$). Table 1 shows that 51% of respondents were females, 53% had some form of high school or higher education; and 53% had gross household incomes below \$25,000 per year in 1991. The majority of respondents were married (64%), resided with at least one other person (67%), and lived in a house (69%). Furthermore, 56% of respondents reported that they were either current or past smokers, and 72% drank alcohol either “sometimes” or “regularly”.

Insert Table 1

Agreement between Survey and Routinely Collected Data

Table 2 shows the observed agreement between the survey and routinely collected data for the health care utilisation measures. The proportion of observed agreement for the contact

utilisation measures (77.1% to 93.6%) was higher than the agreement for the volume utilisation measures (13.6% to 83.5%). According to Landis and Koch's scale (1977), the strengths of observed agreement for contact utilisation measures ranged from substantial to almost perfect compared with poor to moderate strengths of observed agreement for the volume utilisation measures. When κ was calculated, however, poor to substantial levels of agreement were seen for the contact utilisation measures. The substantial level of beyond chance agreement was found for any stay in hospital [$\kappa=0.771$, 95% Confidence interval (CI): 0.718 to 0.824] and any contact with PT or chiropractor ($\kappa=0.685$, 95% CI: 0.622 to 0.748). However, the poor to fair agreement was observed for any contact with a GP ($\kappa=0.195$, 95% CI: 0.126 to 0.265) and any contact with other medical specialists ($\kappa=0.351$, 95% CI: 0.286 to 0.417).

Poor to moderate strengths of beyond chance agreement was shown for the volume utilisation measures. ICC showed moderate agreement for length of stay in hospital (ICC=0.50, 95% CI: 0.45 to 0.55) and number of contacts with PT or chiropractor (ICC=0.41, 95% CI: 0.36 to 0.46); fair agreement was found for the number of contacts with GP (ICC=0.34, 95% CI: 0.25 to 0.42). The poorest agreement was found for the number of contacts with other medical specialists (ICC=0.25, 95% CI: 0.13 to 0.35).

Insert Table 2

Comparison of Contact and Volume Utilisation Measures

The levels of under- and over-reporting between survey and routinely collected data among seniors is shown in Tables 3 and 4. Among the contact utilisation measures, the proportion of

over- and underreporting for any stay in a hospital (3.3% and 3.1% respectively) were relatively small and balanced each other out (Table 3). However, respondents significantly overestimated their contact with a GP, and PT or chiropractor, and underestimated their contact with other medical specialists ($p \leq 0.05$). More specifically, 16.4% of seniors overreported contact with GPs in the past year compared with 4.3% who underreported. For any contact with a PT, 6.0% of seniors overreported visits compared with 2.0% who underreported. As well, 15.6% of seniors underreported visits with other medical specialists, while 7.3% overreported. Therefore, the net reporting error ranged from 4.0% for contact with a PT to 12.1% for contact with a GP.

Among the volume utilisation measures (Table 4), there was no significant difference between the over- and underreporting for length of stay at a hospital and number of contacts with a PT or chiropractor. However, respondents significantly underreported their contact with GPs and other medical specialists (Table 4). Among patients who reported to have at least one contact with their GP in the past year, 9.0% overreported contact by one, 5.0% overreported contact by two, and 12.9% overreported contact by 3 or more. Similarly, 10.3% underreported contact with a GP by one, 8.0% underreported contact by two, and 41.1% underreported contact by three or more. Overall 26.9% of respondents overreported the number of contacts with GPs compared with 59.4% who underreported their contact ($p < 0.001$).

Among respondents who reported to have at least one contact with other medical specialists in the past year, 5.7% overreported contact by one, 2.4% overreported contact by two, and 1.4% overreported contact by three or more. Similarly, 12.0% underreported contact with a medical specialist by one, 9.5% underreported contact by two, and 49.7% underreported contact by three or more. Overall, 9.6% of respondents overestimated the number of contacts with other medical specialists compared with 71.2% who underestimated their contact ($p < 0.001$).

Insert Tables 3 and 4

Logistic and Linear Regression

The logistic regression results for the contact utilisation measures are shown in Table 5. Due to missing data for the independent factors, sample sizes for the logistic regression was decreased by 55 cases for the hospital contact and volume utilisation measures, and 54 cases for the health professional contact and volume utilisation measures. Most models had a good overall goodness-of-fit on Hosmer and Lemeshow statistics. Among statistically significant results, the odds of males agreeing on any stay in a hospital was lower than that of females (Odds Ratio (OR)=0.57, 95% CI: 0.33 to 0.98). The odds of seniors aged 75 years and over agreeing on any stay in a hospital were less than those aged 65-74 years old (OR=0.55, 95% CI: 0.32 to 0.95). Seniors who reported incomes of greater than \$25,000 had a significantly greater odds of agreeing on any contact with both a GP (OR=1.60, 95% CI: 1.14 to 2.24) and other medical specialists (OR=1.40, 95% CI: 1.01 to 1.94) than seniors with incomes of \$25,000 or less. Seniors who reported very good or excellent self-assessed health status had more than twice the odds of agreeing on any stay in hospital (OR=2.51, 95% CI: 1.18 to 5.34) than those who reported poor, fair or good health. As well, seniors with two or more chronic conditions had a lower odds of agreeing on any contact with a PT or chiropractor (OR for 2 chronic conditions=0.13, 95% CI: 0.02 to 0.99 and OR for 3 or more chronic conditions=0.12, 95% CI: 0.02 to 0.91) than seniors who did not report any chronic conditions. However, respondents with three or more chronic conditions had a greater odds of agreeing on any contact with a medical specialist (OR=1.77, 95%

CI:1.02 to 3.05).

The linear regression results for the volume utilisation measures are shown in Table 6. For the length of stay, statistical significance ($\beta=-7.73$, $p\text{-value}<0.0001$) was found only for the self-reported health status. Seniors who had reported a lower health status tended to overreport length of stay in the self-reported data. For number of contacts with GPs, the significant variables were again self reported health status ($\beta=-1.91$, $p\text{-value}<0.0004$) and number of reported chronic conditions ($\beta=0.89$, $p\text{-value}<0.0001$). An increase in chronic condition results in more overreporting. For number of contacts with PTs or chiropractors, significant relationships were found for gender ($\beta=-1.06$, $p\text{-value}<0.0255$) and self-reported health status ($\beta=-1.92$, $p\text{-value}<0.0425$).

Insert Tables 5 and 6

Discussion

The purpose of this study was to examine the level of agreement between self-reported and routinely collected health care utilisation data, as well as to explore the salient factors that may be associated with agreement between the two data sources. Our findings showed substantial to almost perfect observed agreement between survey and routinely collected data for the contact utilisation measures. This supports findings by Glandon and colleagues (1992), who studied a group of 234 adults aged 62 years and older and found that 91.6% of the sample accurately reported contact with a physician within the past 6 months. As well, Wallihan and colleagues

(1999) also reported that 88.4% of adults aged 60 years and over from a clinical study population accurately reported a hospitalisation in the past 12 months.

A wide range of observed agreement, from poor to almost perfect observed agreement, was found for the volume utilisation measures. More specifically, poor observed agreement was found for the number of contacts with a GP and other medical specialists in the past year, and almost perfect observed agreement was found for length of stay in a hospital and for number of contacts with a PT or chiropractor. In the latter case, however, the agreement was mainly the result of a large proportion of respondents reporting no health service use. We can assume that it is easier for respondents to accurately report no use of health services over the past year compared with respondents who had many encounters. When looking only at respondents who had contact with the health care system, observed agreement for the exact length of stay at a hospital was found to be 17.0% (35/206), a value slightly lower than a recently published result of 26.5% by Wallihan and colleagues (1999). Similarly, our observed agreement of 9.8% (96/984) for number of contacts with a GP is lower than the value of 28.2% found by Glandon and colleagues (1992). Lower levels of observed agreement in our findings compared with previous studies might be largely explained by the differences in study populations. Our study utilised community-based non-institutionalised seniors rather than patients who received medical care from a clinic or HMO (Wallihan, Stump and Callahan 1999; Glandon, Counte and Tancredi 1992). Compared with a clinical group, our community sample of seniors may have been more likely to be healthier and less likely to see a physician or to stay overnight at a hospital. Therefore, they may also have been less likely to remember the frequency of their contacts with health professionals compared with clinical seniors patients.

Our study also found that the beyond chance agreement measured by κ for any contact

with a GPs was much lower than expected from our observed agreement. The κ showed poor agreement of 0.195 compared with substantial observed agreement of 79.3%. This paradoxical result may have occurred because high observed values can often be associated with low levels of κ when marginal totals are symmetrically unbalanced in a 2 by 2 table (Feinstein and Cicchetti 1990). In this situation, therefore, the observed agreement is a closer approximation of agreement than κ .

More detailed analyses between survey and administrative data related to the contact utilisation measures showed no difference between the proportion of overestimation and underestimation for any stay at a hospital. However, respondents significantly overestimated any contact with a GP and PT or chiropractor, and significantly underestimated any contact with other medical specialists. Lack of difference between the proportion of over- and underreporting for any stay in a hospital may be because hospitalization is highly salient event and likely to be remembered. Roos and Havens (1991) have also found that more salient events like surgical admissions may be more likely to be remembered than routine medical admissions.

As expected, the volume utilisation measures showed larger disagreement than the contact utilisation measures. No differences were found between the proportions over- and underestimated for length of stay at a hospital and for number of contacts with a PT or chiropractor over the past 12 months. However, respondents significantly underestimated the number of contacts with GPs and other medical specialists. A number of studies have shown that the inaccuracy of recalling an event may be a function of the number of services used. Both Wallihan and colleagues (1999) and Roberts and colleagues (1996) found that the inaccuracy of estimations increased as the volume of services increased. Among respondents who reported at least one hospitalization in the past 12 months, 26.5% were exact in their estimation of length of

stay at a hospital, 43.8% either over- or underestimated their stay by one to three days, and another 29.7% either over- or underestimated their stay by four or more days (Wallihan, Stump and Callahan 1999). Therefore, while individuals may remember whether they visited a doctor or were admitted a hospital, they rarely remember them as countable series (Raina 1995).

Significant disagreements occurred in both contact and volume utilisation despite a previous study by Raina (1995) who found high test-retest reliability related to the any contact and frequency of contact with health professionals. Kappa values (κ) of 0.83 to 0.91 were found for any contact with a GP and other medical specialists, and κ of 0.69 to 0.90 were found for the frequency of contact with a GP and other medical specialists. At least four other reasons may explain the significant over- and underreporting of both contact and volume utilisation measures in our study. Firstly, confusion in the definition of survey questions or terminology may have occurred. Related to the OMH administrative data bases, if a health professional billed for health care services that was not identical to the respondents understanding of the term “contact”, then our analysis would show disagreement. For example, a respondent’s interpretation of contact with a health professional may have included phone calls to the physician’s office, compared with the administrative meaning of contact which is a face to face meeting that results in a bill to the OMH. In addition, there is also a possibility that elderly respondents did not clearly understand the difference between contact with a GP versus other medical specialists in providing primary care services. Although 16.4% of respondents overreported contact in the survey for a GP while a similar number (15.6%) underreported for a medical specialist, further analysis (not shown) did not support the notion that respondents are mistaking specialists for GPs. Respondents who overreported GP contact were actually more likely to overreport, rather than underreport, specialist contact. Therefore respondent, confusion between contact with a GP

and other medical specialists was not evident in our study. From the perspective of the physician, discrepancies may have occurred from inaccurate billing (e.g. specialists providing primary care services and therefore billing as a GP). However, as reported earlier, these discrepancies were minimal since the differences between the two data bases occurred in less than 2% of the listed clinicians.

Secondly, discrepancies between survey and routinely collected data may be due to differences in reporting systems. Related to PT and chiropractic services, in Ontario, Canada, only some physiotherapy or chiropractic clinics are allowed to bill OHIP. All other clinics are privately-owned and bill patients directly. Their data, may not necessarily be collected and included in the administrative data base. As well, if visits to a PT occur under a hospital outpatient physiotherapy department, they would not appear in the routinely collected administrative data but rather under a global hospital budget. Discrepancies related to contact with a GP might result when an individual visits their physician for follow-up care, but the data does not get recorded appropriately. Discrepancies related to reporting of overnight stays at a hospital are influenced by the time an individual was admitted to a hospital. For example, a patient admitted at 3:00 am may believe that he or she is spending that night in the hospital. However, because the admission was missed on a hospital's midnight census, the patient would not be counted as having stayed that night in the administrative data base.

Thirdly, the discrepancy between self-reported and routinely collected data may also be affected by the quality of the recorded health data. Romm & Putnam (1981) compared medical records with verbatim transcripts of outpatient visits and found that only 59% of information present in the transcripts was found in the medical record. Problems related to the recording of medical data include inadequate recording of information by physicians, improper or incomplete

recording of discharge form information, and miscoding of the data (Rawson, Malcolm and D'Arcy 1997; Marrie, Duant and Sealy, 1987; Doremus and Michensi 1983; Demlo, Campbell and Brown 1978). Rawson and D'Arcy (1991) noted that there are many opportunities for biases and errors to occur from first contact of a patient with a health professional to the recording of this event in a data base. The reliability of the information may be affected by potential problems ranging from minor inaccuracies to complete misrepresentation of data (Romm and Putnam 1981).

Fourthly, the accuracy of reporting health care utilisation may be affected by factors that decrease recall memory among seniors. A number of studies in the literature have shown that problems of recall memory increase among older adults who were advanced in age (Weiderholt, Cahn, Butters, et al. 1993; Spiers and Kutsik 1995). Wiederholt and colleagues (1993), studied the performance of neuropsychological tests on 1,692 community-dwelling subjects aged 55 to 94 years and found that the short-term and long-term recall ability decreased progressively with age. In addition, they found that women performed better than men on memory tests even with advancing age. As well, both men and women with some college education performed better on most tests than men and women with high school education (Wiederhoit, Cahn, Butters, et al. 1993). Although recall bias may have played an important role in our results, our sample of older adults were a randomly selected group of community-dwelling seniors who were more likely to be younger in age and have a higher household income than non-respondents. Therefore, our findings may overestimate the proportion of agreement compared with the general population seniors. The length of our recall period, 12 months, may also have had an effect on the respondents' ability to remember the number of contacts they had with health care professionals.

Unlike previous studies in the area, we have not assumed a gold standard for either the

administrative or self-reported data. Our findings suggests that, depending on the source of data, contact utilisation measures may differ by almost one quarter. Disagreement ranged from 6.4% for any stay in a hospital to 22.9% for contact with other medical specialists. Similarly, volume utilisation measures between the two data sources may differ by as much as 85%. Failure to recognize these discrepancies between survey and administrative data may lead to the establishment of inappropriate health care policies.

The logistic regression analyses in our study suggested an important association between socio-demographic and health status variables, and in the level of agreement between survey data and routinely collected data. Agreement generally decreased if respondents were male, aged 75 years and older, had incomes of less than \$25,000, and had a lower (poor/fair/good versus very good/excellent) self-assessed health status. The linear regression suggested that poor or fair health status and male gender tended to be associated with underreporting on volume measures in the surveys. This is consistent with previous reports that also found greater inaccuracies in self-reports occurring in respondents with increasing age and disability, less social supports, less education, and lower income (Wallihan, Stump and Callahan 1999; Gladdon, Counte and Tancredi 1992).

These large differences in estimates suggest that previous studies, which relied only on one source of data to estimate health care use, may be seriously biased in one direction or another. Therefore, investigators seeking to study the future use of health care services among community dwelling older adults can not solely rely on self-reported data or routinely collected data. Self-reported and administrative data are considered useful sources, however investigators need to confront the validity and quality of their data when attempting to interpret findings. Future studies should examine the magnitude of error in measuring the contact and volume of

health services utilisation, as well as other factors that may more accurately predict health care use among community-based seniors using these data sources.

References

Branch L., A. Jette, and C. Evashwick. 1981. "Towards understanding elders' health service utilisation." *J Community Health* 7: 80-92.

Bush T.L., S.R. Miller, A.L. Golden, and W.E. Hale. 1989. "Self-report and medical record report agreement of selected medical conditions in the elderly." *Am J Public Health* 79: 1554-6.

Chappell N, L. Strain, and A. Blandford. 1986. *Aging and health care: A social perspective*. Toronto: Holt, Rinehart and Winston of Canada, Limited.

Cummings S.R., M.C. Nevitt, and S. Kidd. 1988. "Forgetting falls: the limited accuracy of recall of falls in the elderly." *J Am Geriatr Soc* 36: 613-16.

Demlo L.K., P.M. Campbell, and S.S. Brown. 1978. "Reliability of information abstracted from patient's medical records." *Med Care* 16: 995-1005.

Doremus H.D. and E.M. Michensi. 1983. "Data quality. An illustration of its potential impact upon a diagnosis-related group's case mix index and reimbursement." *Med Care* 21: 1001-11.

Feinstein A.R. and D.V. Cicchetti. 1990. "High agreement but low kappa: I. The problems of two paradoxes." *J Clin Epidemiol* 43: 543-49.

Fleiss J.L. 1986. *The design and analysis of clinical experiments*. New York: John Wiley and Sons.

Glandon G.L., M.A. Counte, and D. Tancredi. 1992. "An analysis of physician utilisation by elderly persons: Systematic differences between self-report and archival information." *J Gerontol* 47(5): S245-S252.

Holland C.A. and P.M. Rabbitt. 1991. "Ageing memory: use versus impairment." *Br J Psych* 82(Pt 1): 29-38.

Landis J.R. and G.G. Koch. 1977. "The measurement of observer agreement for categorical data." *Biometrics* 33: 159-74.

Mackenbach J.P., C.W.N. Looman, and J.B.W. Van der Meer. 1996. "Differences in the misreporting of chronic conditions, by level of education: The effect on inequalities in prevalence rates." *Am J Public Health* 86: 706-11.

Marrie T.J., H. Durant, and E. Sealy. 1987. "Pneumonia. The Quality of Medical Records Data." *Med Care* 25: 20-4.

Mossey J.M., B. Havens, and F.D. Wolinsky. 1989. "The consistency of formal health care utilisation: physician and hospital utilisation." In *Aging and Health Care*, edited by M.G. Ory and K. Bond, pp. 81-98. London: Routledge Press.

Muller C. 1986. "Review of twenty years of research on medical care utilisation." *Health Services Research* 21: 129-44.

National Center for Health Statistics. 1987. "Reporting chronic conditions in the National Health Interview Survey: A review of tendencies from evaluation studies and methodological tests." *Vital Health Statistics*: 105-13.

Ory M.G. and K. Bond. 1989. "Introduction: health care for an aging population." In *Aging and Health Care*, edited by M.G. Ory and K. Bond, pp. 1-21. London: Routledge Press.

Prigerson H.G., P.K. Maciejewski, and R.A. Rosenheck. 1999. "The effects of marital dissolution and marital quality on health and health service use among women." *Med Care* 37(9): 858-73.

Raina P., B. Bonnett, D. Waltner-Toews, C. Woodward, and T. Abernathy. 1999. "How reliable are selected scales from population-based health surveys? An analysis among seniors." *Can J Public Health* 90(1): 60-4.

Raina P, D. Waltner-Toews, B. Bonnett, C. Woodward, and T. Abernathy. 1999. "Influence of companion animals on the physical and psychological health of older people: An analysis of a one-year longitudinal study. *J Am Geriatr Soc* 47: 323-29.

Raina P. 1995. *The impact of pet ownership and attachment on the health and health care use of the elderly in Wellington County, Ontario* [dissertation]. Guelph, Ontario: University of Guelph,.

Rawson N.S.B., E. Malcolm, and C. D'Arcy. 1997. "Reliability of the recording of schizophrenia and depressive disorder in the Saskatchewan health care datafiles." *Soc Psych Psych Epidemiol* 32: 191-199.

Roberts R.O., E.J. Bergstralh, L. Schmidt, and S.J. Jacobsen. 1996. "Comparison of self-reported and medical record health care utilisation measures." *J Clin Epidemiol* 49: 989-95.

Romm F.J., and S.M. Putnam. 1981. "The validity of the medical record." *Med Care* 14: 310-15.

Roos N., and B. Havens. 1991. "Predictors of successful aging: A twelve-year study of Manitoba elderly." *Am J Public Health* 81, no.1: 63-68.

Spiers M.V., and D.M. Kutsik. 1995. "Self-reported memory of medication use by the elderly." *Am J Health-System Pharm* 52(9):985-90.

Statistics Canada. 1994. *National Population Health Survey 1994-1995*. Ottawa: Statistics Canada.

Strout P.E., and J.L. Fleiss. 1979. "Intraclass correlations: Uses in assessing rater reliability." *Psychol Bulletin* 86: 420-28.

Wallihan D.B., T.E. Stump, and C.M. Callahan. 1999. "Accuracy of self-reported health services use and patterns of care among urban older adults." *Med Care* 37(7): 662-70.

Wan T.T.H. 1989. "The behavioral model of health care utilisation by older people." In *Aging and Health Care*, edited by M.G. Ory and K. Bond, pp. 52-71. London: Routledge Press.

Wiederholt W.C., D Cahn, N.M. Butters, D.P. Salmon, D. Krits-Silverstein, and E. Barrett-Connor. 1993. "Effects of age, gender and education on selected neuropsychological tests in an elderly community cohort." *J Am Geriatr Soc* 41(6): 639-47.

Table 1: Distribution of demographic characteristics of non-institutionalised community dwelling seniors aged 65 years and older (n=1038), 1992

Variables	n (%)
<u>Gender</u>	
Female	529 (51)
Male	509 (49)
<u>Age Groups</u>	
65-69	353 (34)
70-74	332 (32)
75-79	187 (18)
80 & over	166 (16)
<u>Household Income*</u>	
Under \$11,999	119 (11)
\$12,000-\$24,999	441 (42)
\$25,000-49,999	331 (32)
\$50,000 & over	103 (10)
<u>Education</u>	
Below high school	488 (47)
High school and above	550 (53)
<u>Marital Status</u>	
Married	664 (64)
Widowed	291 (28)
Other (e.g., never married/ Divorced/separated)	83 (8)
<u>Living Arrangement</u>	
Alone	343 (33)
Living with someone	695 (67)
<u>Type of dwelling</u>	
House	716 (69)
Apartment	197 (19)
Other (e.g., trailer, townhouse and other types of dwellings)	125 (12)
<u>Smoking</u>	
Current smokers	104 (10)
Past smokers	477 (46)
Never smokers	455 (44)
<u>Alcohol†</u>	
Never drinkers	280 (27)
Sometimes drinkers	466 (45)
Regular drinkers	290 (28)

* Missing information accounted for the remaining 5% of the data.

† Status of alcohol drinking: never drinkers (did not drink alcohol at all), sometime drinkers (drank 1 to 3 times a month or less than once a month), and regular drinkers (drank at least once a week)

Table 2: Summary of exact agreement between survey and routinely collected data on health care utilisation measures among seniors aged 65 years and older, 1992

Contact Utilisation Measure (n=sample size)	Observed Agreement: n (%)	Kappa (95% Confidence Intervals)
Any stay in hospital (n=1036)	970 (93.6)	0.771 (0.718, 0.824)
Any contact with general practitioner (n=1028)	815 (79.3)	0.195 (0.126, 0.264)
Any contact with physiotherapist or chiropractor (n=1028)	945 (91.9)	0.685 (0.622, 0.748)
Any contact with other medical specialists (n=1028)	793 (77.1)	0.351 (0.286, 0.417)

Volume Utilisation Measure (n=sample size)	Observed Agreement: n (%)	Intraclass Correlation Coefficient (95% Confidence Intervals)
Length of stay in hospital (n=1036)	865 (83.5)	0.50 (0.45, 0.55)
Number of contacts with general practitioner (n=1028)	140 (13.6)	0.34 (0.25, 0.42)
Number of contacts with physiotherapist or chiropractor (n=1028)	844 (82.1)	0.41 (0.36, 0.46)
Number of contacts with other medical specialists (n=1028)	197 (19.2)	0.25 (0.13, 0.35)

Table 3: Comparison between survey responses and routinely collected data on the contact utilisation measures among seniors aged 65 years and older, 1992

Contact Utilisation Measure (n=sample size)	Survey Response*	Data base Indication*	Frequency	Percent
Any stay in hospital (n=1036)				
Agreement	no	no	830	80.1
	yes	yes	140	13.5
Disagreement: Overreporting in survey	yes	no	34	3.3
	Underreporting in survey	no	yes	32
Any contact with general practitioner (n=1028)				
Agreement	no	no	44	4.3
	yes	yes	771	75.0
Disagreement: Overreporting in survey	yes	no	169	16.4**
	Underreporting in survey	no	yes	44
Any contact with physiotherapist or chiropractor (n=1028)				
Agreement	no	no	832	80.9
	yes	yes	113	11.0
Disagreement: Overreporting in survey	yes	no	62	6.0**
	Underreporting in survey	no	yes	21
Any contact with other medical specialists (n=1028)				
Agreement	no	no	114	11.0
	yes	yes	679	66.1
Disagreement: Overreporting in survey	yes	no	75	7.3**
	Underreporting in survey	no	yes	160

* "No" indicates no stay or contact reported in the data source, "yes" indicates that at least one stay or contact was reported in the data source

** Indicates a significant difference between the two types of disagreement ($p \leq 0.05$)

Table 4: Comparison between survey responses and administrative data base on the volume utilisation measures among seniors aged 65 years and older,1992

	Length of stay in hospital (n=1036)		Number of contacts with general practitioner (n=1028)		Number of contacts with physiotherapist or chiropractor (n=1028)		Number of contacts with other medical specialists (n=1028)	
	Freq	%	Freq	%	Freq	%	Freq	%
Agreement*								
On responses of zero	830	80.1	44	4.3	832	80.9	114	11.1
On responses of one or more	35	3.4	96	9.3	12	1.2	83	8.1
Disagreement								
Over-reporting in survey by 1	22	2.1	93	9.0	17	1.6	59	5.7
Over-reporting in survey by 2	10	1.0	51	5.0	9	0.9	25	2.4
Over-reporting in survey by 3 or more	47	4.5	133	12.9	75	7.3	15	1.4
TOTAL OVER-REPORTING	79	7.6	277	26.9**	101	9.8	182	9.6**
Under-reporting in survey by 1	31	3.0	106	10.3	14	1.4	123	12.0
Under-reporting in survey by 2	15	1.4	82	8.0	11	1.1	98	9.5
Under-reporting in survey by 3 or more	46	4.4	423	41.1	58	5.6	511	49.7
TOTAL UNDER-REPORTING	92	8.9	611	59.4**	83	8.1	732	71.2**

* Agreement of the volume utilisation measures included zero responses (e.g. respondents reported zero for length of stays at a hospital and the data base was also zero) and responses of one or more (e.g. respondents of the survey reported one or more nights of stay at a hospital and database also reported one or more nights stay)

** Indicates a significant difference between the two types of disagreement (p<0.0010)

Table 5: Logistic regression results for agreement/disagreement* on the contact utilisation measures among seniors aged 65 years and older, 1992

Independent Factors**	Odds Ratio (95% Confidence Interval)			
	Any stay in hospital (n=981)	Any contact with GP (n=974)	Any Contact with PT or chiropractor (n=974)	Any Contact with other medical specialists (n=974)
Gender:				
Female	Ref	Ref	Ref	Ref
Male	0.57 (0.33, 0.98)	0.81 (0.58, 1.11)	1.20 (0.74, 1.93)	1.05 (0.77, 1.45)
Age:				
65-74 years	Ref	Ref	Ref	Ref
75+ years	0.55 (0.32, 0.95)	1.20 (0.85, 1.72)	1.00 (0.61, 1.64),	0.81 (0.59, 1.12)
Income:				
≤\$25,000	Ref	Ref	Ref	Ref
>\$25,000	1.60 (0.89, 2.87)	1.60 (1.14, 2.24)	1.12 (0.68, 1.83)	1.40 (1.01, 1.94)
Self-Assessed Health Status:				
Poor, fair or good	Ref	Ref	Ref	Ref
Very good or excellent	2.51 (1.18, 5.34)	0.84 (0.41, 1.70)	1.55 (0.71, 3.41)	0.68 (0.34, 1.36)
Self-Assessed Emotional Health Status:				
Poor, fair or good	Ref	Ref	Ref	Ref
Very good or excellent	2.11 (0.94, 4.73)	0.68 (0.30, 1.53)	0.58 (0.19, 1.73)	0.94 (0.46, 1.91)
Number of chronic Conditions:				
0	Ref	Ref	Ref	Ref
1	0.43 (0.05, 3.71)	0.78 (0.42, 1.45)	0.43 (0.05, 3.74)	1.32 (0.72, 2.43)
2	0.19 (0.03, 1.49)	1.19 (0.64, 2.21)	0.13 (0.02, 0.99)	1.60 (0.88, 2.90)
3 or more	0.22 (0.03, 1.65)	1.54 (0.86, 2.74)	0.12 (0.02, 0.91)	1.77 (1.02, 3.05)
Hosmer and Lemeshow Goodness-of-Fit	7.85 (df=8) sig=0.4481	14.08 (df=8) sig=0.0798	8.39 (df=8) sig=0.3965	6.50 (df=8) sig=0.5911

* Agreement/disagreement includes all responses from survey and administrative data (e.g., agreement="yes/yes" and "no/no" and disagreement="no/yes" and "yes/no").

** The variables education, marital status, living arrangement, type of dwelling, and current status of smoking and alcohol drinking were omitted from the analyses because they were not significantly associated with agreement between the two sources of data ($p > 0.05$) and did not contribute to the overall model.

Table 6: Linear regression for predictors of differences between self-reported and MOH data for volume measures among seniors aged 65 years and older, 1992

Independent Factors	Dependent Variables											
	Length of stay in hospital (n=981)			# of contacts with GP (n=974)			# of contacts with PT (n=974)			# of contacts with medical specialists (MS) (n=974)		
	β	SE	P-Value	β	SE	P-Value	β	SE	P-Value	β	SE	P-Value
Constant	6.49	2.13	0.0024	4.44	0.90	0.0001	2.57	1.52	0.0467	0.59	0.72	0.4152
Gender (0=Female & 1=Male)	-0.80	0.65	0.2247	-0.05	0.27	0.8130	-1.06	0.47	0.0255	-0.16	0.24	0.4960
Age (0=65-74 & 1=75+)	-1.08	0.69	0.1196	-0.27	0.29	0.3572	-0.24	0.50	0.6368	0.41	0.25	0.1068
Income(0= \leq 25,000 & 1= $>$ 25,000)	0.76	0.67	0.2588	-0.43	0.28	0.1244	-0.74	0.48	0.1288	0.21	0.24	0.3853
Self-Assessed Health (0=poor/good & 1=Very good/excellent)	-7.73	1.31	0.0001	-1.99	0.55	0.0004	-1.92	0.95	0.0425	-0.22	0.48	0.6415
Self-Assessed Emotional Health (0=poor/good & 1=Very good/excellent)	1.91	1.43	0.1824	-1.07	0.60	0.0741	0.20	1.04	0.8413	-0.44	0.52	0.3947
# of Chronic Conditions	0.43	0.33	0.2044	0.89	0.14	0.0001	0.46	0.24	0.0572	0.22	0.12	0.0762
Length of Stay in MOH data	-0.54	0.03	0.0001		---			---			---	
# of contacts with GP in MOH data		---		-0.80	0.01	0.0001		---			---	
# of contacts with PT in MOH data		---			---		-0.16	0.49	0.0011		---	
# of contacts with MS in MOH data		---			---			---		-0.84	0.10	0.0001

** The variables education, marital status, living arrangement, type of dwelling, and current status of smoking and alcohol drinking were omitted from the analyses since they were not significantly associated with agreement between the two sources of data ($p > 0.05$) and did not contribute to the overall model.

SEDAP RESEARCH PAPERS

Number	Title	Author(s)
No. 1:	Population Aging and Its Economic Costs: A Survey of the Issues and Evidence	F.T. Denton B.G. Spencer
No. 2:	How Much Help Is Exchanged in Families? Towards an Understanding of Discrepant Research Findings	C.J. Rosenthal L.O. Stone
No. 3:	Did Tax Flattening Affect RRSP Contributions?	M.R. Veall
No. 4:	Families as Care-Providers Versus Care-Managers? Gender and Type of Care in a Sample of Employed Canadians	C.J. Rosenthal A. Martin-Matthews
No. 5:	Alternatives for Raising Living Standards	W. Scarth
No. 6:	Transitions to Retirement: Determinants of Age of Social Security Take Up	E. Tompa
No. 7:	Health and Individual and Community Characteristics: A Research Protocol	F. Béland S. Birch G. Stoddart
No. 8:	Disability Related Sources of Income and Expenses: An Examination Among the Elderly in Canada	P. Raina S. Dukeshire M. Denton L.W. Chambers A. Scanlan A. Gafni S. French A. Joshi C. Rosenthal
No. 9:	The Impact of Rising 401(k) Pension Coverage on Future Pension Income	W.E. Even D.A. Macpherson
No. 10:	Income Inequality as a Canadian Cohort Ages: An Analysis of the Later Life Course	S.G. Prus
No. 11:	Are Theories of Aging Important? Models and Explanations in Gerontology at the Turn of the Century	V.L. Bengtson C.J. Rice M.L. Johnson
No. 12:	Generational Equity and the Reformulation of Retirement	M.L. Johnson
No. 13:	Long-term Care in Turmoil	M.L. Johnson L. Cullen D. Patsios
No. 14:	The Effects of Population Ageing on the Canadian Health Care System	M.W. Rosenberg

SEDAP RESEARCH PAPERS

Number	Title	Author(s)
No. 15:	Projections of the Population and Labour Force to 2046: Canada	F.T. Denton C.H. Feaver B.G. Spencer
No. 16:	Projections of the Population and Labour Force to 2046: The Provinces and Territories	F.T. Denton C.H. Feaver B.G. Spencer
No. 17:	Location of Adult Children as an Attraction for Black and White Elderly Migrants in the United States	K.-L. Liaw W.H. Frey J.-P. Lin
No. 18:	The Nature of Support from Adult <i>Sansei</i> (Third Generation) Children to Older <i>Nisei</i> (Second Generation) Parents in Japanese Canadian Families	K.M. Kobayashi
No. 19:	The Effects of Drug Subsidies on Out-of-Pocket Prescription Drug Expenditures by Seniors: Regional Evidence from Canada	T.F. Crossley P. Grootendorst S. Korkmaz M.R. Veall
No. 20:	Describing Disability among High and Low Income Status Older Adults in Canada	P. Raina M. Wong L.W. Chambers M. Denton A. Gafni
No. 21:	Parental Illness and the Labour Supply of Adult Children	P.T.Léger
No. 22:	Some Demographic Consequences of Revising the Definition of 'Old' to Reflect Future Changes in Life Table Probabilities	F.T. Denton B.G. Spencer
No. 23:	Geographic Dimensions of Aging: The Canadian Experience 1991-1996	E.G. Moore D. McGuinness M.A. Pacey M.W. Rosenberg
No. 24:	The Correlation Between Husband's and Wife's Education: Canada, 1971-1996	L. Magee J. Burbidge L. Robb
No. 25:	The Effect of Marginal Tax Rates on Taxable Income: A Panel Study of the 1988 Tax Flattening in Canada	M.-A. Sillamaa M.R. Veall
No. 26:	The Stability of Self Assessed Health Status	T.F. Crossley S. Kennedy

SEDAP RESEARCH PAPERS

Number	Title	Author(s)
No. 27:	How Do Contribution Limits Affect Contributions to Tax-Preferred Savings Accounts?	K. Milligan
No. 28:	The Life Cycle Model of Consumption and Saving	M. Browning T.F. Crossley
No. 29:	Population Change and the Requirements for Physicians: The Case of Ontario	F.T. Denton A. Gafni B.G. Spencer
No. 30:	Nonparametric Identification of Latent Competing Risks and Roy Duration Models	G. Colby P. Rilstone
No. 31:	Simplified Estimation of Multivariate Duration Models with Unobserved Heterogeneity	G. Colby P. Rilstone
No. 32:	Structural Estimation of Psychiatric Hospital Stays	G. Colby P. Rilstone
No. 33:	Have 401(k)s Raised Household Saving? Evidence from the Health and Retirement Study	G.V. Engelhardt
No. 34:	Health and Residential Mobility in Later Life: A New Analytical Technique to Address an Old Problem	L.M. Hayward
No. 35:	2 ½ Proposals to Save Social Security	D. Fretz M.R. Veall
No. 36:	The Consequences of Caregiving: Does Employment Make a Difference	C.L. Kemp C.J. Rosenthal
No. 37:	Fraud in Ethnocultural Seniors' Communities	P.J.D. Donahue
No. 38:	Social-psychological and Structural Factors Influencing the Experience of Chronic Disease: A Focus on Individuals with Severe Arthritis	P.J. Ballantyne G.A. Hawker D. Radoeva
No. 39:	The Extended Self: Illness Experiences of Older Married Arthritis Sufferers	P.J. Ballantyne G.A. Hawker D. Radoeva
No. 40:	A Comparison of Alternative Methods to Model Endogeneity in Count Models. An Application to the Demand for Health Care and Health Insurance Choice	M. Schellhorn
No. 41:	Wealth Accumulation of US Households: What Do We Learn from the SIPP Data?	V. Hildebrand

SEDAP RESEARCH PAPERS

Number	Title	Author(s)
No. 42:	Pension Portability and Labour Mobility in the United States. New Evidence from SIPP Data.	V. Andrietti V. Hildebrand
No. 43:	Exploring the Effects of Population Change on the Costs of Physician Services	F.T. Denton A. Gafni B.G. Spencer
No. 44:	Reflexive Planning for Later Life: A Conceptual Model and Evidence from Canada	M.A. Denton S. French A. Gafni A. Joshi C. Rosenthal S. Webb
No. 45:	Time Series Properties and Stochastic Forecasts: Some Econometrics of Mortality from the Canadian Laboratory	F.T. Denton C.H. Feaver B.G. Spencer
No. 46:	Linear Public Goods Experiments: A Meta-Analysis	J. Zelmer
No. 47:	Local Planning for an Aging Population in Ontario: Two Case Studies	L.M. Hayward
No. 48:	Management Experience and Diversity in an Ageing Organisation: A Microsimulation Analysis	T. Wannell M. Gravel
No. 49:	Resilience Indicators of Post Retirement Well-Being	E. Marziali P. Donahue
No. 50:	Continuity or Change? Older People in Three Urban Areas	J. Phillips M. Bernard C. Phillipson J. Ogg
No. 51:	Intracohort Income Status Maintenance: An Analysis of the Later Life Course	S.G. Prus
No. 52:	Tax-Preferred Savings Accounts and Marginal Tax Rates: Evidence on RRSP Participation	K. Milligan
No. 53:	Cohort Survival Analysis is Not Enough: Why Local Planners Need to Know More About the Residential Mobility of the Elderly	L.M. Hayward N.M. Lazarowich
No. 54:	Unemployment and Health: Contextual Level Influences on the Production of Health in Populations	F. Béland S. Birch G. Stoddart

SEDAP RESEARCH PAPERS

Number	Title	Author(s)
No. 55:	The Timing and Duration of Women's Life Course Events: A Study of Mothers With At Least Two Children	K.M. Kobayashi A. Martin-Matthews C.J. Rosenthal S. Matthews
No. 56:	Age-Gapped and Age-Condensed Lineages: Patterns of Intergenerational Age Structure Among Canadian Families	A. Martin-Matthews K. M. Kobayashi C.L. Rosenthal S.H. Matthews
No. 57:	The Relationship between Age, Socio-Economic Status, and Health among Adult Canadians	S.G. Prus
No. 58:	Measuring Differences in the Effect of Social Resource Factors on the Health of Elderly Canadian Men and Women	S.G. Prus E. Gee
No. 59:	APOCALYPSE NO: Population Aging and the Future of Health Care Systems	R.G. Evans K.M. McGrail S.G. Morgan M.L. Barer C. Hertzman
No. 60:	The Education Premium in Canada and the United States	J.B. Burbidge L. Magee A.L. Robb
No. 61:	Student Enrolment and Faculty Recruitment in Ontario: The Double Cohort, the Baby Boom Echo, and the Aging of University Faculty	B.G. Spencer
No. 62:	The Social and Demographic Contours of Contemporary Grandparenthood: Mapping Patterns in Canada and the United States	C.L. Kemp
No. 63:	Changing Income Inequality and the Elderly in Canada 1991-1996: Provincial Metropolitan and Local Dimensions	E.G. Moore M.A. Pacey
No. 64:	Mid-life Patterns and the Residential Mobility of Older Men	L.M. Hayward
No. 65:	The Retirement Incentive Effects of Canada's Income Security Programs	M. Baker J. Gruber K. Milligan
No. 66:	The Economic Well-Being of Older Women Who Become Divorced or Separated in Mid and Later Life	S. Davies M. Denton

SEDAP RESEARCH PAPERS

Number	Title	Author(s)
No. 67:	Alternative Pasts, Possible Futures: A “What If” Study of the Effects of Fertility on the Canadian Population and Labour Force	F.T. Denton C.H. Feaver B.G. Spencer
No. 68:	Baby-Boom Aging and Average Living Standards	W. Scarth M. Souare
No. 69:	The Invisible Retirement of Women	L. McDonald
No. 70:	The Impact of Reference Pricing of Cardiovascular Drugs on Health Care Costs and Health Outcomes: Evidence from British Columbia – Volume I: Summary	P.V. Grootendorst L.R. Dolovich A.M. Holbrook A.R. Levy B.J. O'Brien
No. 71:	The Impact of Reference Pricing of Cardiovascular Drugs on Health Care Costs and Health Outcomes: Evidence from British Columbia – Volume II: Technical Report	P.V. Grootendorst L.R. Dolovich A.M. Holbrook A.R. Levy B.J. O'Brien
No. 72:	The Impact of Reference Pricing of Cardiovascular Drugs on Health Care Costs and Health Outcomes: Evidence from British Columbia – Volume III: ACE and CCB Literature Review	L.R. Dolovich A.M. Holbrook M. Woodruff
No. 73:	Do Drug Plans Matter? Effects of Drug Plan Eligibility on Drug Use Among the Elderly, Social Assistance Recipients and the General Population	P. Grootendorst M. Levine
No. 74:	Living Alone and Living with Children: The Living Arrangements of Canadian and Chinese-Canadian Seniors	M.A. Pacey
No. 75:	Student Enrolment and Faculty Recruitment in Ontario: The Double Cohort, the Baby Boom Echo, and the Aging of University Faculty (Revised and updated version of No. 61)	B.G. Spencer
No. 76:	Gender Differences in the Influence of Economic, Lifestyle, and Psychosocial Factors on Later-life Health	S.G. Prus E. Gee
No. 77:	Asking Consumption Questions in General Purpose Surveys	M. Browning T.F. Crossley G. Weber
No. 78:	A Longitudinal Study of the Residential Mobility of the Elderly in Canada	Y. Ostrovsky
No. 79:	Health Care in Rural Communities: Exploring the Development of Informal and Voluntary Care	M.W. Skinner M.W. Rosenberg

SEDAP RESEARCH PAPERS

Number	Title	Author(s)
No. 80:	Does Cognitive Status Modify the Relationship Between Education and Mortality? Evidence from the Canadian Study of Health and Aging	J.C. Brehaut P. Raina J. Lindsay
No. 81:	Agreement between Self-Reported and Routinely Collected Health Care Utilisation Data among Seniors	P. Raina V. Torrance-Rynard M. Wong C. Woodward